

HOSTED BY




ELSEVIER

Contents lists available at ScienceDirect

China University of Geosciences (Beijing)

Geoscience Frontiers

journal homepage: www.elsevier.com/locate/gsf

Research paper

The inception of a Paleotethyan magmatic arc in Iberia

M.F. Pereira^{a,*}, A. Castro^b, C. Fernández^c^aIDL, Departamento de Geociências, ECT, Universidade de Évora, Apartado 94, 7001-554 Évora, Portugal^bDepartamento de Geología, Universidad de Huelva, Spain^cDepartamento de Geodinámica y Paleontología, Universidad de Huelva, Spain

ARTICLE INFO

Article history:

Received 17 July 2013

Received in revised form

24 January 2014

Accepted 12 February 2014

Available online 12 March 2014

Keywords:

Permo-Carboniferous

Calc-alkaline arc-related batholiths

Pangaea

Paleotethys subduction

Variscan and Cimmerian cycles

ABSTRACT

This paper presents a compilation of recent U–Pb (zircon) ages of late Carboniferous–early Permian (LC–EP) calc-alkaline batholiths from Iberia, together with a petrogenetic interpretation of magma generation based on comparisons with Mesozoic and Tertiary Cordilleran batholiths and experimental melts. Zircon U–Pb ages distributed over the range ca. 315–280 Ma, indicate a linkage between calc-alkaline magmatism, Iberian orocline generation and Paleotethys subduction. It is also shown that Iberian LC–EP calc-alkaline batholiths present unequivocal subduction-related features comparable with typical Cordilleran batholiths of the Pacific Americas active margin, although geochemical features were partially obscured by local modifications of magmas at the level of emplacement by country rock assimilation. When and how LC–EP calc-alkaline batholiths formed in Iberia is then discussed, and a new and somewhat controversial interpretation for their sources and tectonic setting (plume-assisted relamination) is suggested. The batholiths are proposed to have formed during the subduction of the Paleotethys oceanic plate (Pangaea self-subduction) and, consequently, they are unrelated to Variscan collision. The origin of the Iberian batholiths is related to the Eurasian active margin and probably represents the inception of a Paleotethyan arc in the core of Pangaea.

© 2015, China University of Geosciences (Beijing) and Peking University. Production and hosting by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

There is currently general consensus regarding global palaeogeographic views of Pangaea during late Carboniferous–early Permian times (LC–EP; ca. 315–280 Ma), which place Iberia at the core of this supercontinent, near the Eurasian active margin located to the east (Cocks and Torsvik, 2006; Stampfli and Kozur, 2006, and references therein). Surprisingly, in spite of this widely-accepted palaeogeographic proximity, the likely relationship between the source of Iberian LC–EP calc-alkaline magmas and the subduction of the Paleotethys oceanic plate has never been explored.

This point of view was first discussed in a general way for the Alpine arc (Finger and Steyrer, 1990) with regard to the Variscan granitoids of the Tauern area. But the existence of a wide magmatic province in the Alpine domain (Catalonia, Pyrenees, Calabria,

Sardinia, Corsica, Provence, Tuscan nappes, Briançonnais domain, the southern and eastern Alps), characterized by the transition from LC–EP calc-alkaline plutonism to early to mid-Permian post-collisional or extension related magmatism was documented by Stampfli (1996) and references therein. The inception of the subduction of Paleotethys in the Permian and a representation of subduction-related magmatic belts were later presented by Stampfli (2001).

Until now, LC–EP calc-alkaline magmas in Iberia have been linked by a number of researchers with a range of sources and tectonic settings associated with Variscan collision. Among these, are: (1) the reworking of different crustal protoliths, including metaigneous and metasedimentary rocks (Bea et al., 2003); (2) the reworking of oceanic metabasic rocks accreted to mid-to-lower crustal levels (Villasca et al., 2009); (3) the involvement of mantle-derived melts or meta-igneous lower crust (Fernández Suárez et al., 2011); and (4) a primarily igneous rock source stemming from depleted mantle or meta-igneous lower crust and derived by partial melting of heterogeneous metasedimentary detrital rocks in the mid-crust (Neiva et al., 2009). Recently, a new tectonic model was presented for the origin of LC–EP calc-alkaline

* Corresponding author. Tel.: +351 266 745301; fax: +351 266 745397.
E-mail address: mpereira@uevora.pt (M.F. Pereira).

Peer-review under responsibility of China University of Geosciences (Beijing).

magmatism in Iberia (Gutiérrez-Alonso et al., 2008, 2011; Fernández Suárez et al., 2011; Weil et al., 2012). According to this interpretation, mantle and lower-mid crust melting was triggered by lithosphere delamination coeval with the formation of the Iberian orocline. The proposed tectonic model suggests that closure of the Paleotethys Ocean was probably accommodated by unique Pangaea rotation and self-subduction during LC–EP times (e.g., Gutiérrez-Alonso et al., 2008). This permits a possible relationship between the development of the Eurasian active margin and magmatism occurring simultaneously in Iberia (Gutiérrez-Alonso et al., 2011).

Although many authors consider the LC–EP calc-alkaline magmatism to show geochemical characteristics demonstrably similar to magmas formed in subduction settings, this alternative interpretation of magma sources has never been considered in detail previously. Traditionally, the consensus has been that Iberian LC–EP calc-alkaline magmas are associated with the Variscan orogenic cycle (“late Variscan batholiths”), despite their being ca. 60–80 Ma younger than the subduction of the Rheic Ocean (ca. 385–365 Ma; Martínez-Catalan et al., 2007, 2009; Nance et al., 2010). A combination of factors have, in our view, precluded the satisfactory clarification of the sources and tectonic setting of ca. 315–280 Ma calc-alkaline magmatism in Iberia: (1) the scarcity of precise absolute ages for accurately determining the relationship between different calc-alkaline batholiths and the timing of deformation; (2) the lack of a robust petrogenetic model for explaining the origin of magmas and comparing them with other extensively studied examples of arc-related batholiths and corresponding petrologic and experimental constraints; and (3) the general acceptance of a single, consensual interpretation that supports the development of this calc-alkaline magmatic suite within the framework of the Variscan orogenic cycle. This paradigm disregards the wider context of the complex formation of Pangaea and the spatial proximity of Iberia relative to the Eurasian active margin during the LC–EP closing of the Paleotethys Ocean.

In this paper, an alternative interpretation is proposed for the source of the Iberian LC–EP calc-alkaline granitoids and mafic rocks within a subduction setting. In support of this new and controversial interpretation are the following data sets: (1) a compilation of recently published U–Pb (zircon) geochronology data obtained for the Iberian LC–EP calc-alkaline magmatism (Neiva et al., 2009; Fernández-Suárez et al., 2011; Díaz Alvarado et al., 2013; Pereira et al., 2014); and (2) a comparison of major compositional relations between the Iberian batholiths and the Cordilleran batholiths of North Patagonia (Castro et al., 2011) and those of eastern Peninsular Ranges (Lee et al., 2007), and with experimental residues and melts using a new robust model for simulating the generation of new crust in arcs (Castro et al., 2013). We conclude by examining the origin of ca. 315–280 Ma calc-alkaline batholiths in Iberia in relation to the Eurasian active margin and discuss whether the batholiths provide evidence for the inception of a Paleotethyan arc in the core of Pangaea.


2. Calc-alkaline Iberian batholiths: U–Pb zircon ages and compositions

The late Palaeozoic closure of the Rheic Ocean, following the collision of Gondwana and Laurussia, contributed to a mountain-building event during the formation of Pangaea (Variscan Orogen; Martínez-Catalan et al., 2007, 2009). Iberia, located in the western branch of the European Variscan belt, includes a ca. 1000 km-long magmatic province with a huge volume of LC–EP calc-alkaline batholiths extending from the Central-Iberian Zone to the southern Pyrenees (Fig. 1a). This calc-alkaline magmatic suite is dominated by I-type granodiorites with subordinate mafic rocks and S-type

granites generated by crustal anatexis (Castro et al., 2002, 2003; Bea et al., 2003; Neiva et al., 2009; Villaseca et al., 2009; Fernández-Suárez et al., 2011; Díaz Alvarado et al., 2013). These calc-alkaline batholiths intruded Neoproterozoic to early Palaeozoic sedimentary successions that are dominated by detrital rocks with associated minor igneous bodies.

A compilation of recently published U–Pb geochronology data from Iberian LC–EP batholiths is presented in Table 1. The U–Pb (zircon) data show that calc-alkaline magmatism occurred in Iberia over an extended period of ca. 35 million years (LC–EP; Table 1; Fig. 1b). In Iberia, three main phases of Variscan deformation are described, which enable different magmatic suites to be distinguished (e.g., Castro et al., 2002; Dias et al., 2002; Martínez-Catalan et al., 2007; Fernández-Suárez et al., 2011; Gutiérrez-Alonso et al., 2011; Díaz Alvarado et al., 2013). In accordance with this classification, LC–EP magmatic suites have been described as syn-D₃ and post-D₃ (“late Variscan”) because they show sharp intrusive contacts with previous Variscan structures (D₁ and D₂ deformation phases). D₁ structures are related to the Variscan collision and the emplacement of allochthonous units (ca. 365–345 Ma; Escuder Viruete et al., 1994; Dallmeyer et al., 1997; Rubio Pascual et al., 2013), while D₂ structures are linked to Variscan extensional collapse and widespread crustal anatexis (ca. 345–320 Ma; Escuder Viruete et al., 1994; Dallmeyer et al., 1997; Martínez Catalán et al., 2004; Castiñeiras et al., 2008; Pereira et al., 2012a,b; Rubio Pascual et al., 2013), after cessation of Rheic subduction. Lower Carboniferous (LC) calc-alkaline intrusive bodies coeval with D₂ have been interpreted as coming from a mantle-derived magma that interacted, in different degrees, with crustal melts (Dias et al., 1998; Montero et al., 2004; Romeo et al., 2006; Moita et al., 2009; Pereira et al., 2009; Lima et al., 2011). Younger Variscan structures (D₃ deformation phase) are associated with the formation of the Iberian orocline (ca. 320–300 Ma; Gutiérrez-Alonso et al., 2008, 2011; Pereira et al., 2012a).

The geochemical composition of Iberian LC–EP batholiths is summarized in Fig. 2. The ca. 315–280 Ma magmatic series from Iberia are divided into (Castro et al., 2002, and references therein): (1) a silicic (SiO₂ > 53 wt.%) calc-alkaline (I-type) series forming large batholiths with predominantly granodiorite composition; (2) a mafic rock association (diorite-gabbro) that occurs both in peribatholithic massifs near the main contacts of the large granodiorite intrusions and intruding into migmatitic complexes; and (3) a peraluminous leucogranite (S-type) series, commonly associated with *in situ* migmatization. Fig. 2 includes geochemical diagrams (major elements) showing essential variations of representative Iberian LC–EP granitoids and mafic rocks (ca. 315–280 Ma). Iberian LC–EP granitoids are well grouped along curved trends characteristic of Cordilleran batholiths (Fig. 2a). Mafic rocks are partly cumulates formed as part of the calc-alkaline differentiation trend, and are partly derived from wet basaltic magmas (appinites) that are commonly associated with the Iberian batholiths, and show characteristic and unequivocal arc geochemical signatures (Castro et al., 2003) (Fig. 2a). The variation of the molar Mg[#] shows a plateau region for rocks within the silica range 60–70 wt.%, which is a characteristic feature of Cordilleran granitoid batholiths (for example, Patagonian batholiths: Pankhurst et al., 1999; Hervé et al., 2007; Castro et al., 2011; and Sierra Nevada and Peninsular Ranges batholiths: Lee et al., 2007) (Fig. 2b). The continuous linear increase of the molar K ratio with decreasing MgO is also characteristic of calc-alkaline magma fractionation from a parental diorite precursor (Fig. 2c). An increase in aluminosity (A parameter: de la Roche, 1978; Debon and Le Fort, 1983) with fractionation is a characteristic feature of calc-alkaline (cafemic) trends (Fig. 2d). In all diagrams shown, the effects of the assimilation of local country rock (pelitic



to the low pressure and moderate water cotectic liquids (thick dashed line: [Sisson et al., 2005](#); [Castro, 2013](#)). Several samples tend to deviate from the cotectic line due to the local effects of pelitic rock assimilation ([Díaz Alvarado et al., 2011](#)).

Table 1
Compilation of LC–EP calc-alkaline Iberian batholith ages (ca. 315–280 Ma).

Rocks	Location	Age (Ma)	Method	Author
Bt granodiorites	CIZ (Gredos)	312	U/Pb zrn, SHRIMP	Díaz Alvarado et al., 2013
Granodiorites	CIZ (Castelo Branco)	309	U/Pb zrn ID-TIMS	Antunes et al., 2010
Tonalites, gabbros	CIZ (Arges, Toledo, Guajaraz)	308–311	U/Pb zrn, Cameca IMS270	Bea et al., 2006
Bt ± Amph granodiorites	CIZ (Los Pedroches)	308	U/Pb zrn ID-TIMS	Carracedo et al., 2009
Peraluminous granites	CIZ (Casal Vasco, Junqueira)	308–311	U/Pb zrn, mnz ID-TIMS	Valle Aguado et al., 2005
Monzogranites	CIZ (Cota)	307	U/Pb zrn, mnz ID-TIMS	Valle Aguado et al., 2005
Bt granites and Crd granites	CIZ (Béjar)	307	U/Pb zrn, SHRIMP	Zeck et al., 2007
Granitoids	CIZ (Gredos)	307	U/Pb zrn, SHRIMP	Díaz Alvarado et al., 2013
Bt monzogranites	CIZ (Berlengas island)	307	U/Pb zrn ID-TIMS	Valverde-Vaquero et al., 2011
Monzogranite	CIZ-OMZ (Nisa)	307–309	U/Pb zrn, SHRIMP	Solá et al., 2009
Granites	CIZ (Gouveia)	307–310	U/Pb zrn ID-TIMS	Neiva et al., 2012
Mafic and ultramafic rocks	CIZ (Gredos)	307–322	²⁰⁷ Pb/ ²⁰⁶ Pb zrn, ID-TIMS	Montero et al., 2004
Bt monzogranites	CIZ (Braga-Celeirós)	306–311	U/Pb zrn, mnz ID-TIMS	Dias et al., 1998
Mafic rocks	CIZ (Toledo, La Bastida)	306–311	Cameca IMS270	Bea et al., 2006
Tonalite	CIZ-OMZ (Aldeia da Mata)	306	U/Pb zrn, SHRIMP	Solá et al., 2009
Gabbro and norites	CIZ (Béjar)	306	U/Pb zrn, SHRIMP	Zeck et al., 2007
Leucogranites	CIZ (Gredos)	305	U/Pb zrn, SHRIMP	Díaz-Alvarado et al., 2013
Leucogranites	CIZ (Cerro Mogábar)	304	U/Pb zrn, ID-TIMS	Carracedo et al., 2008
Crd monzogranite	CIZ (Gredos)	304	U/Pb zrn, SHRIMP	Díaz Alvarado et al., 2013
Migmatitic hornfelses	CIZ (Gredos)	303	U/Pb zrn, SHRIMP	Díaz Alvarado et al., 2013
Granites and leucogranites	CIZ (La Cabrera)	302	²⁰⁷ Pb/ ²⁰⁶ Pb zrn, ID-TIMS	Casquet et al., 2004
Two-mica monzogranites	CIZ (Gouveia)	302	U-Th-Pb mnz, SHRIMP	Neiva et al., 2009
Leucogranites	CIZ (Briteiros)	300	U/Pb zrn, mnz ID-TIMS	Dias et al., 1998
Bt monzogranites	CIZ (Vila Pouca de Aguiar)	299	U/Pb zrn, ID-TIMS	Martins et al., 2009
Quartz diorite	E-Pyrenees (Cap Creus-Tudela)	299	U/Pb zrn, SHRIMP	Druguet et al., 2014
Granodiorites	S-Pyrenees (Boí, Montellá)	299–301	U/Pb zrn, SHRIMP	Pereira et al., 2014
Cold-granitoids	CIZ	299	U/Pb zrn, LA-ICPMS	Gutierrez-Alonso et al., 2011
Granites	CIZ (Gouveia)	297	U/Pb zrn ID-TIMS	Neiva et al., 2012
Granites	CIZ (Gouveia)	297–304	U/Pb mnz ID-TIMS	Neiva et al., 2012
Hot-granitoids	CIZ-CZ-WALZ	292–309	U/Pb zrn, LA-ICPMS/ID-TIMS	Gutierrez-Alonso et al., 2011 and references therein
Two-mica monzogranite	CIZ (La Cabrera)	292	U/Pb zrn, ID-TIMS	Valverde-Vaquero et al., 1997
Granodiorite	E-Pyrenees (Roses)	291	U/Pb zrc, SHRIMP	Druguet et al., 2014
Bt granites	CIZ (Gerês-Peneda)	290–296	U/Pb zrn, mnz, ID-TIMS	Dias et al., 1998
Leucogranite	CIZ (Gouveiras)	289	U-Th-Pb mnz, SHRIMP	Neiva et al., 2009
Cold-granitoids	CIZ-WALZ	286–306	U/Pb zrn, LA-ICPMS/ID-TIMS	Gutierrez-Alonso et al., 2011 and references therein
Granodiorites	S-Pyrenees (Vielha)	276	U/Pb zrn, SHRIMP	Pereira et al., 2014

3. Discussion

The generation of Iberian LC–EP calc-alkaline batholiths in a tectonic setting distinct from subduction, conflicts with essential thermal requirements and phase equilibria constraints on magma generation and fractionation. Moreover, the generation of these large batholiths in response to a supposed late-Variscan extension is incompatible with the overall calc-alkaline affinity of the magmas.

It is broadly recognized that granitic rocks of extensional (e.g., back-arc) settings are of alkaline affinity (A-type), with strong Fe enrichment, high Na/K ratios and typically meta-aluminous features (Eby, 1990). The chemical composition of the Iberian LC–EP calc-alkaline batholiths, however, is distinct from that of the Permian to Cretaceous NE China batholiths with granitic rocks of A-type and transitional I-type to A-type affinities that formed in one of the world's most outstanding areas of lithospheric extension (Wu et al., 2002, 2005).

The Iberian LC–EP calc-alkaline batholiths also display transitions to locally-formed anatectic granites (S-type). It is known that the greater or lesser presence of associated S-type granites largely depends on the level of emplacement and the more or less fertile composition of the crustal lithologies hosting the calc-alkaline batholiths. For example, S-types granites are scarce or absent in large regions of the Patagonian batholiths in South America where rocks hosting the intrusions are mostly volcanics. However, S-type granites are present, and transitions from I-type to S-type are common in other Cordilleran-type batholiths, for example the pre-collisional S-type granites of the Lhasa terrane as part of the calc-

alkaline Gangdese batholith (Zhang et al., 2013), and the anatectic granites from the Famatina arc system associated with the Gondwana active margin in Argentina (Grosse et al., 2011). In the case of Iberia, the emplacement of the LC–EP calc-alkaline batholiths into a fertile middle crustal level dominated by a several kilometer-thick sequence of Neoproterozoic greywackes is the reason for extensive anatexis over large regions close to the intrusion contacts. Migmatites are common at the contacts, with anatectic granites showing the same age as the intruding magma (e.g., Refugio del Rey leucogranite and Gredos granodiorites; Díaz Alvarado et al., 2013). The interaction between the intrusive granodiorites and the fertile metasedimentary host gave rise to mechanical interactions between the magma and partially molten host, leading to local hybridization of the intrusive magma, which becomes more peraluminous and more potassic. Criteria for identifying these local processes have been determined by means of combined geochemical and experimental studies (Díaz Alvarado et al., 2011).

The ultimate origin of the granodiorite magmas in relation to subduction is a problem of general scope that has been widely discussed in previous papers. The lower crust is often cited as the source of calc-alkaline batholiths, either for those closely related to recognizable active margins, or those apparently unrelated to coeval subduction.

The lower crust interpretation is based on two important observations: (1) many batholiths are intrusive into the middle crust at pressures ranging from 4 to 8 kbar; and (2) melts produced from the mantle underlying the continental crust must be basaltic. A two-step process consisting of the melting of mantle-derived basaltic rocks previously underplated in the lower crust has been

proposed (Petford and Atherton, 1996; Petford and Gallagher, 2001; Hawkesworth and Kemp, 2006). Plume-assisted relamination is a new concept in subduction-related magmatism emerging from a variety of independent approaches: (1) thermomechanical numerical experiments (Gerya et al., 2004; Gerya and Meilick, 2011; Vogt et al., 2012); (2) experimental phase equilibria of batholith-forming magmas (Castro et al., 2010; Castro, 2013, 2014); and (3) mass balance calculations (Hacker et al., 2011). Essentially, the concept involves the generation of magmas of intermediate composition (andesite-diorite) by the melting of subducted materials in silicic composite plumes (composed of oceanic crust and sediments) which are finally relaminated at a level below the lower crust, where they split by means of melt segregation into liquids that are emplaced into the middle and upper crust (batholiths) and residues that remain in the lower crust (mafic granulites). Relaminated andesite magmas may reach the lower continental crust at high temperatures of about 1000 to 1100 °C, containing a crystal proportion of about 50% according to predictions from phase equilibrium experiments. Temperatures of the same order (ca. 1000 °C) are recorded by lower crust granulite xenoliths around the world (for example, Paso de Indios xenoliths; Castro et al., 2011, 2013, and references therein). Furthermore, the characteristic low-water content of calc-alkaline batholiths, with an initial water content of about 1–2 wt.% H₂O, is compatible with high temperatures of about 1000 °C at the time of magma segregation from composite underplated plumes.

In the case of the Iberian LC–EP calc-alkaline granitoids, which have been considered to be unrelated to coeval subduction (the Rheic Ocean had closed ca. 60–80 Ma earlier; e.g., Martínez-Catalán et al., 2007, 2009; Pereira et al., 2012b), the lower crust is often cited as their source. In our view, several factors indicate an extra-crustal origin for the Iberian LC–EP calc-alkaline magmas, suggesting a link to a subduction setting similar to that of the Cordilleran batholiths formed during the Mesozoic and Tertiary at the Pacific active margin of the Americas (e.g., Sierra Nevada: Lee et al., 2006, 2007, and Patagonia: Pankhurst et al., 1999; Parada et al., 1999; Herve et al., 2007; Castro et al., 2011). A detailed discussion of the arguments supporting an off-crust generation of calc-alkaline batholiths is provided in other papers (e.g., Castro et al., 2010, 2013; Castro, 2014).

We have shown that the I-type granodiorites forming the Iberian LC–EP batholiths have the features of calc-alkaline silicic magmatism characteristic of active continental margins like the Andes (Patagonian batholith) and the North American Cordillera (Sierra Nevada batholith). Another unequivocal sign of a subduction setting is the occurrence of mafic rocks (appinites). These represent calc-alkaline mafic systems with typical arc signatures (Murphy, 2013). They resemble the geochemical signature of high-Mg andesites (HMA) and sanukitoids, formed by the reaction of subduction-derived melts with the overlying mantle (Kelemen et al., 2003). However, the possibility that they were alkaline hydrous magmas enriched with crustal components by contamination from country rocks has been suggested for particular cases (Scarrow et al., 2009). The problem here relates to classical discussion on the origin of the crustal signatures of arc magmas as acquired by either: (1) the assimilation of lower crust materials by basaltic magmas (Hildreth and Moorbath, 1988); or (2) the incorporation of crustal components (including sedimentary rocks) into the mantle source of magmas through subduction erosion (Stern, 1991). The former (crustal contamination) may be disregarded on the basis of simple mass balance calculations because the preserved basaltic nature of magmas would have been destroyed by silica enrichment associated with assimilation or magma mixing (Castro et al., 2003). The latter implies that crustal signatures originated in a metasomatized mantle source or through the reaction between

melts from the subduction zone and the overlying peridotite mantle (Kelemen et al., 2003). The relationships between these subduction-related mafic rocks and large Iberian LC–EP calc-alkaline batholiths are crucial for providing an understanding of their origins. I-type granodiorites represent silicic magmas, possibly derived by fractionation from a diorite (andesite) precursor that could not have been generated by melting the lower crust, or through the melting of a metasomatized mantle. It has recently been proposed that the origin of these subduction-related mafic rocks is associated with plume-assisted relamination (Vogt et al., 2012; Castro et al., 2013) as part of a new paradigm for magma generation in the supra-subduction mantle wedge (Gerya et al., 2004).

We realize, however, that the onset of magmatism related to Paleotethys subduction (Eo-Cimmerian cycle) may have overlapped late-magmatic Variscan events (Druguet et al., 2014; Pereira et al., 2014). It is possible that some Iberian LC–EP magmatic events took place in the transition from the end of Variscan collision to the continuing subduction of the Paleotethys active margin. With the available chronological and geochemical data it is difficult to unequivocally trace the boundary between these two magmatic events of different origin. A detailed geochronology and geochemistry study of the wet-basaltic (normally shoshonitic) magmatism that is present in Iberia may reveal important clues on this problem.

In summary, the use of a robust petrogenetic model for explaining the origin of magmas, and a comparison with other extensively studied examples of arc-related batholiths, was essential for deciphering the subduction-related geochemical signatures of the Iberian magmas that were partially obscured by local modifications of the magmas at the level of emplacement. The presence of several km-thick sequences of Ediacaran and Cambrian detrital Al-rich sedimentary host rocks explains the widespread geochemical modifications of the intruding magmas, which underwent local assimilation and conversion to hybrid cordierite monzogranites. The late generation of S-type peraluminous leucogranites is related to the local thermal effects of the intruding batholiths.

The compilation of recent geochronology U–Pb (zircon) data provides conclusive evidence that precludes a genetic connection between Iberian LC–EP calc-alkaline magmatism and Variscan collision. The period during which calc-alkaline magmatism was generated (ca. 315–280 Ma) coincides with the development of the Iberian orocline (Gutiérrez-Alonso et al., 2008; Fernández-Suárez et al., 2011; Weil et al., 2012; Johnston et al., 2013). Therefore, orocline formation and Pangaea self-subduction (Gutiérrez-Alonso et al., 2008, 2011) acted together to cause the generation of the LC–EP calc-alkaline magmas of Iberia in a setting similar to that of typical Cordilleran batholiths (Fig. 3).

In palaeogeographic reconstructions of Pangaea formation during the LC–EP, Iberia is bounded in the west by the Variscan suture and in the east by the Paleotethys Ocean (Cocks and Torsvik, 2006; Stampfli and Kozur, 2006). We suggest that Paleotethys subduction that led to the collision between the Cimmerian terranes (detached from the non-collisional northern margin of Gondwana) and the Eurasian active margin was responsible for the inception of a Paleotethyan magmatic arc in Iberia. We note that the generation of arc-related calc-alkaline magmas in Iberia accounts for a possible link between the Variscan and Cimmerian cycles (Druguet et al., 2014; Pereira et al., 2014). The early stages of Paleotethys Ocean closure were followed by the generation of calc-alkaline arc-related magmas in Iberia, simultaneously with lithosphere-scale bending that produced the accommodation of large continental blocks with associated ductile faulting and the resulting emplacement of large batholiths (Gutiérrez-Alonso et al.,




Figure 2. Geochemical diagrams (major elements) showing essential variations of representative granitoids of the Iberian Massif (<315 Ma). Based on geochemical data from Casillas (1989); Ugidos et al. (1997); Bea et al. (1999); López-Plaza and López-Moro (2003); Antunes (2006); Antunes et al. (2008); Orejana et al. (2009) and Díaz Alvarado et al. (2011). Granitoids forming these large Iberian batholiths, classically considered to be late Variscan-late granodiorites of Capdevila et al. (1973) and Castro et al. (2002), are well grouped along curved trends characteristic of Cordilleran batholiths. (a) The CaO-MgO variation diagram can be taken as a proxy of cotectic calc-alkaline evolution trends typically found in arc magmas. The curved trend defines a line of cotectic liquids (LCC) in agreement with experimental melt compositions in calc-alkaline systems. A typical experimental line is traced in the diagrams (dark red dashed line) according to data in Castro (2013). Some of the mafic rocks are cumulates belonging to the calc-alkaline differentiation trend. Other mafic rocks are derived from wet basaltic magmas (appinites), which are commonly associated with the Iberian LC-EP calc-alkaline batholiths and show characteristic and unequivocal arc




Figure 3. Geodynamic model of the inception of a Paleotethyan arc in Iberia (Adapted from [Castro et al., 2013](#); [Johnston et al., 2013](#); [Pereira et al., 2014](#) and references therein). (A, B, C1) Sequential block diagrams showing the subduction of the Paleotethys oceanic lithosphere (dark-blue), development of the Iberian orocline, the Paleotethyan arc and retroarc (pink), and related strike-slip tectonics. White arrow = approximate direction of convergence between the Paleotethys oceanic lithosphere and the southern limit of the Variscan realm (black lines represent the trace of Variscan structures). Green arrow = lithosphere delamination. Black arrow = slab rollback of the oceanic plate. Grey arrow = orogen-normal principal compressive stress responsible for orocline development. (C2) Block diagrams showing the oceanic-continental subduction zone. Plumes (orange) represent the result of dehydration of the subducted crust, aqueous fluid transport, partial melting, melt extraction and melt emplacement in the form of extrusive volcanics and intrusive plutons (red).

geochemical signatures ([Castro et al., 2003](#)). (b) Variation of the molar $Mg^\#$ shows a plateau region for rocks within the silica range 60–70 wt.%, which is a characteristic feature of Cordilleran granitoid batholiths (e.g., Patagonian batholith: [Pankhurst et al., 1999](#); [Herve et al., 2007](#); [Castro et al., 2011](#); and Sierra Nevada and Peninsular Ranges batholiths: [Lee et al., 2006, 2007](#)). (c) Linear and continuous increase of the molar K ratio with decreasing MgO is characteristic of calc-alkaline magma fractionation from a parental diorite precursor. (d) The increase in aluminosity (A parameter: [de la Roche, 1978](#); [Debon and Le Fort, 1983](#)) with fractionation is characteristic of calc-alkaline (cafemic) trends. In all diagrams, the effects of assimilation of local country rock (pelitic metasediments) are apparent. Contaminated samples depart from the cotectic trends and point to the composition of pelitic migmatites. (e) Pseudoternary projection into the compositional space $F(Fe + Mg + Mn)$ -Anorthite(An)-Orthoclase(Or). This diagram is used to discriminate between processes of restite unmixing and peritectic restite entrainment that operate in S-type granites ([Stevens et al., 2007](#); [Clemens and Stevens, 2012](#)), giving rise to geochemical trends linking anatectic melts and restites (labelled granite-nebulite trends), from processes related to cotectic magma crystallization and fractionation that characterize granodiorite-tonalite batholiths. Most rocks of the calc-alkaline series in the Iberian batholiths plot close to experimental cotectic lines, preferentially close to the low pressure and moderate water cotectic liquids (thick dashed line: [Sisson et al., 2005](#); references for other cotectic lines, red, green and black in [Castro, 2013](#)). Several samples tend to deviate from the cotectic line due to local effects of pelitic rock assimilation ([Díaz Alvarado et al., 2011](#)). Grey areas in (e) represent the fields of S- and I-type granitoids. Large star represents the probable composition of the parental magma precursor to batholiths ([Castro et al., 2013](#)). Characteristic minerals are plotted for reference. Mineral abbreviations after [Kretz \(1983\)](#).

2011; Weil et al., 2012). We propose that the formation of the Iberian LC–EP calc-alkaline batholiths constitutes the earliest magmatic event of the Eurasian active margin. The northward subduction of Paleotethys was responsible for the widespread late Carboniferous calc-alkaline intrusions and volcanism found in the Variscan Alpine–Mediterranean domain (Cortesogno et al., 1998; Stampfli and Borel, 2002). The gradual spread of the Neotethys Ocean and the northward drift of the Cimmerian terranes against the Eurasian active margin during the Permian may have caused the eastward migration of Paleotethys subduction and the associated magmatic arc during the course of the Permian. We further suggest that slab rollback of the Paleotethys oceanic plate during the Permian and Triassic caused the opening of back-arc basins within the Eurasian margin from Austria to China. Later, these back-arc basins were gradually closed during the late Triassic to late Jurassic Cimmerian orogenic cycle (Stampfli and Borel, 2002).

4. Conclusions

LC–EP calc-alkaline batholiths (ca. 315–280 Ma) which formed in Iberia several tens of millions of years from the end of the Rheic Ocean subduction and the time of Gondwana and Laurussia collision (ca. 360 Ma) are unrelated to Variscan orogeny. The magmatism preserves unequivocal subduction-related features, comparable with typical Cordilleran batholiths of the Pacific American active margin. It is proposed that Iberian LC–EP calc-alkaline batholiths were formed during the subduction of the Paleotethys oceanic plate (Pangaea self-subduction) representing the inception of a Paleotethyan arc related to the Eurasian active margin.

Acknowledgements

The authors acknowledge the financial support that comes from the Research Projects “GOLD” (FCT-PTDC/GEO-GEO/2446/2012-FCOMP-01-0124-FEDER-029192) and “PLUVOLC” (No. CGL2010-22022). This is a contribution to IGCP projects 597 – Amalgamation and Breakup of Pangaea: the type example of the supercontinent cycle and 574-Bending and Bent orogens, and continental ribbons.

References

- Antunes, A., Santos, J.F., Azevedo, M.R., Mendes, M.H., Ribeiro, S., 2010. New petrographical, geochemical and geochronological data for the Reguengos de Monsaraz pluton (Ossa–Morena Zone, SW Iberian Massif, Portugal). *Estudios Geológicos* 66 (1), 25–34.
- Antunes, I., 2006. Mineralogia, Petrologia e Geoquímica de Rocas Granitoides da área de Castelo Branco – Idanha-a-Nova. Unpublished PhD thesis. Universidade de Coimbra, p. 453.
- Antunes, I.M.H.R., Neiva, A.M.R., Silva, M.M.V.G., Corfu, F., 2008. Geochemistry of S-type granitic rocks from the reversely zoned Castelo Branco pluton (central Portugal). *Lithos* 103, 445–465.
- Bea, F., Montero, P., González Lodeiro, F., Talavera, C., Molina, J.F., Scarrow, J.H., Whitehouse, M.J., Zinger, T., 2006. Zircon thermometry and U–Pb ion-microprobe dating of the gabbros and associated migmatites of the Variscan Toledo anatectic complex, Central Iberia. *Journal of the Geological Society of London* 163, 847–855.
- Bea, F., Montero, P., Zinger, T., 2003. The nature, origin, and thermal influence of the granite source layer of Central Iberia. *Journal of Geology* 111, 579–595.
- Bea, F., Montero, P., Molina, J.F., 1999. Mafic precursors, peraluminous granitoids, and late lamprophyres in the Avila batholith: a model for the generation of Variscan batholiths in Iberia. *Journal of Geology* 107, 399–419.
- Capdevila, R., Corretgé, G., Floor, P., 1973. Les granitoides varisques de la Meseta Ibérique. *Bulletin de la Société Géologique de France* 15, 209–228.
- Carracedo, M., Paquette, J.L., Alonso Olazabal, A., Santos Zalduegui, J.F., García de Madinabeitia, S., Tiepolo, M., Gil Ibarra, J.I., 2009. U–Pb dating of granodiorite and granite units of the Los Pedroches batholith. Implications for geodynamic models of the southern Central Iberian Zone (Iberian Massif). *International Journal of Earth Sciences* 98 (7), 1609–1624.
- Casillas, R., 1989. Las asociaciones plutónicas tardihercínicas del Sector Occidental de la Sierra de Guadarrama – Sistema Central Español (Las Navas del Marqués – S. Martín de Valdeiglesias). Petrología, Geoquímica, Génesis y Evolución. Unpublished PhD thesis. Universidad de Madrid, p. 316.
- Casquet, C., Montero, P., Galindo, C., Bea, F., Lozano, R., 2004. Geocronología 207Pb/206Pb en cristal único de circon y Rb–Sr del plutón de La Cabrera (Sierra del Guadarrama). *Geogaceta* 35, 71–74.
- Castiñeiras, P., Villaseca, C., Barbero, L., Martín Romero, C., 2008. SHRIMP U–Pb zircon dating of anatexis in high-grade migmatite complexes of central Spain: implications in the Hercynian evolution of Central Iberia. *International Journal of Earth Sciences* 97, 35–50.
- Castro, A., 2014. The off-crust origin of granite batholiths. *Geoscience Frontiers* 5, 63–75.
- Castro, A., 2013. Tonalite-granodiorite suites as cotectic systems: a review of experimental studies with applications to granitoid petrogenesis. *Earth Science Reviews* 124, 68–95.
- Castro, A., Vogt, K., Gerya, T.V., 2013. Generation of new continental crust by sub-lithospheric silicic-magma relamination in arcs: a test of Taylor's andesite model. *Gondwana Research* 23, 68–95.
- Castro, A., Moreno-Ventas, I., Fernández, C., Vujovich, G., Gallastegui, G., Heredia, N., Martino, R.D., Becchio, R., Corretgé, L.G., Díaz-Alvarado, J., García-Arias, M., Liu, D.-Y., 2011. Petrology and SHRIMP U–Pb zircon geochronology of Cordilleran granitoids of the Bariloche area, Argentina. *Journal of South American Earth Sciences* 32, 508–530.
- Castro, A., Gerya, T., García-Casco, A., Fernández, C., Díaz Alvarado, J., Moreno-Ventas, I., Loew, I., 2010. Melting relations of MORB-sediment mélanges in underplated mantle wedge plumes. Implications for the origin of cordilleran-type batholiths. *Journal of Petrology* 51, 1267–1295.
- Castro, A., Corretgé, L.G., De La Rosa, J.D., Fernández, C., López, S., Chacón, H., 2003. The appinite–migmatite complex of Sanabria, NW Iberian Massif, Spain. *Journal of Petrology* 44, 1309–1334.
- Castro, A., Corretgé, L.G., De La Rosa, J., Enrique, P., Martínez, F.J., Pascual, E., Lago, M., Arranz, E., Galé, C., Fernández, C., Donaire, T., López, S., 2002. Paleozoic magmatism. In: Gibbons, W., Moreno, M.T. (Eds.), *The Geology of Spain*. Geological Society, London, pp. 117–153.
- Clemens, J.D., Stevens, G., 2012. What controls chemical variation in granitic magmas? *Lithos* 134–135, 317–329.
- Cocks, L.R.M., Torsvik, T.H., 2006. European geography in a global context from the Vendian to the end of the Palaeozoic. In: Gee, D.G., Stephenson, R.A. (Eds.), *European Lithosphere Dynamics*. Geological Society, London, pp. 83–95.
- Cortesogno, L., Cassinini, G., Dallagiovanna, G., Gaggero, L., Oggiano, G., Ronchi, A., Seno, S., Vanossi, M., 1998. The Variscan post-collisional volcanism in late Carboniferous–Permian sequences of Ligurian Alps, southern Alps and Sardinia (Italy): a synthesis. *Lithos* 45, 305–328.
- Dallmeyer, R.D., Martínez Catalán, J.R., Arenas, R., Gil Ibarra, J.I., Gutiérrez-Alonso, G., Fariás, P., Aller, J., Bastida, F., 1997. Diachronous Variscan tectono-thermal activity in the NW Iberian Massif: evidence from ⁴⁰Ar/³⁹Ar dating of regional fabrics. *Tectonophysics* 277, 307–337.
- Debon, F., Le Fort, P., 1983. A chemical-mineralogical classification of common plutonic rocks and associations. *Transactions of the Royal Society of Edinburgh: Earth Sciences* 73, 135–149.
- de la Roche, H., 1978. La chimie des roches présentée et interprétée d'après la structure de leur faciès minéral dans l'espace des variables chimiques: fonctions spécifiques et diagrammes qui s'en déduisent. Application aux roches ignées. *Chemical Geology* 21, 63–87 (in French with English abstract).
- Dias, G., Simões, P.P., Ferreira, N., Leterrier, J., 2002. Mantle and crustal sources in the genesis of late-Hercynian granitoids (NW Portugal): geochemical and Sr–Nd isotopic constraints. *Gondwana Research* 5, 287–305.
- Dias, G., Leterrier, J., Mendes, A., Simões, P.P., Bertrand, J.M., 1998. U–Pb zircon and monazite geochronology of post-collisional Hercynian granitoids from the Central Iberian Zone (Northern Portugal). *Lithos* 45, 349–369.
- Díaz Alvarado, J., Fernández, C., Castro, A., Moreno-Ventas, I., 2013. SHRIMP U–Pb zircon geochronology and thermal modeling of multilayer granitoid intrusions. Implications for the building and thermal evolution of the Central System batholith, Iberian Massif, Spain. *Lithos* 175–176, 104–123.
- Díaz Alvarado, J., Castro, A., Fernández, C., Moreno-Ventas, I., 2011. Assessing bulk assimilation in cordierite-bearing granitoids from the Central System batholith, Spain: experimental, geochemical and geochronological constraints. *Journal of Petrology* 52, 223–256.
- Drugué, E., Castro, A., Chichorro, M., Pereira, M.F., Fernández, C., 2014. Zircon geochronology of intrusive rocks from Cap de Creus, Eastern Pyrenees. *Geological Magazine* 151 (6), 1095–1114.
- Eby, G.N., 1990. The A-type granitoids: a review of their occurrence and chemical characteristics and speculations on their petrogenesis. *Lithos* 26, 115–134.
- Escuder Viruete, J., Arenas, R., Martínez Catalán, J.R., 1994. Tectonothermal evolution associated with Variscan crustal extension in the Tormes Gneiss Dome (NW Salamanca, Iberian Massif, Spain). *Tectonophysics* 238, 1–22.
- Faure, M., Cocherie, A., Mézème, A.B., Charles, N., Rossi, Ph., 2010. Middle carboniferous crustal melting in the Variscan Belt: new insights from U–Th–Pb. Monazite and U–Pb zircon ages of the Montagne Noire Axial zone (southern French Massif Central). *Gondwana Research* 18, 653–673.
- Fernández-Suárez, J., Gutiérrez-Alonso, G., Johnston, S.T., Jeffries, T.E., Pastor-Galan, D., Jenner, G.A., Murphy, J.B., 2011. Iberian late-Variscan granitoids: some considerations on crustal sources and the significance of “mantle extraction ages”. *Lithos* 123, 121–132.

- Finger, F., Steyrer, H.P., 1990. I-type granitoids as indicators of a late Paleozoic convergent ocean/continent margin along the southern flank of the Central European Variscan Orogen. *Geology* 18, 1207–1210.
- Gerya, T.V., Meilick, F.I., 2011. Geodynamic regimes of subduction under an active margin: effects of rheological weakening by fluids and melts. *Journal of Metamorphic Geology* 29, 7–31.
- Gerya, T.V., Yuen, D.A., Sevre, E.O.D., 2004. Dynamical causes for incipient magma chambers above slabs. *Geology* 32, 89–92.
- Grosse, P., Bellos, L.I., de los Hoyos, C.R., Larrovere, M.A., Rossi, J.N., Toselli, A.J., 2011. Across-arc variation of the Famatinian magmatic arc (NW Argentina) exemplified by I- and transitional I/S-type Early Ordovician granitoids of the Sierra de Velasco. *Journal of South American Earth Sciences* 32, 110–126.
- Gutiérrez-Alonso, G., Fernández-Suárez, J., Jeffries, T.E., Johnston, S.T., Pastor-Galán, D., Murphy, J.B., Piedad Franco, M., Gonzalo, J.C., 2011. Diachronous post-orogenic magmatism within a developing orocline in Iberia. *Tectonics* 30, TC5008. <http://dx.doi.org/10.1029/2010TC002845>.
- Gutiérrez-Alonso, G., Fernández-Suárez, J., Weil, A.B., Murphy, J.B., Nance, R.D., Corfu, F., Johnston, S.T., 2008. Self-subduction of the Pangaea global plate. *Nature Geoscience* 1 (8), 549–553.
- Hacker, B.R., Kelemen, P.B., Behn, M.D., 2011. Differentiation of the continental crust by remelting. *Earth and Planetary Science Letters* 307, 501–516.
- Hawkesworth, C.J., Kemp, A.I.S., 2006. Evolution of the continental crust. *Nature* 443, 811–817.
- Herve, F., Pankhurst, R.J., Fanning, C.M., Calderon, M., Yaxley, G.M., 2007. The South Patagonian batholith: 150 my of granite magmatism on a plate margin. *Lithos* 97, 373–394.
- Hildreth, W., Moorbath, S., 1988. Crustal contributions to arc magmatism in the Andes of Central Chile. *Contributions to Mineralogy and Petrology* 98, 455–489.
- Johnston, S.T., Weil, A.B., Gutiérrez-Alonso, G., 2013. Oroclines: thick and thin. *GSA Bulletin* 125 (5–6), 643–663.
- Kelemen, P.B., Hanghøj, K., Greene, A., 2003. One view of the geochemistry of subduction-related magmatic arcs, with an emphasis on primitive andesite and lower crust. In: Holland, H.D., Turekian, K.K., Rudnick, R.L. (Eds.), *The Crust: Treatise on Geochemistry*, vol. 3. Elsevier Pergamon, Oxford, pp. 593–659.
- Kretz, R., 1983. Symbols for rock-forming minerals. *American Mineralogist* 68, 277–279.
- Kroner, U., Romer, R.L., 2013. Two plates – many subduction zones: the Variscan orogeny reconsidered. *Gondwana Research* 24, 298–329.
- Lee, C.-T.A., Morton, D.M., Kistler, R.W., Baird, A.K., 2007. Petrology and tectonics of Phanerozoic continent formation: from island arcs to accretion and continental arc magmatism. *Earth and Planetary Science Letters* 263, 370–387.
- Lee, C.T.A., Cheng, X., Horodyskyj, U., 2006. The development and refinement of continental arcs by primary basaltic magmatism, garnet pyroxenite accumulation, basaltic recharge and delamination: insights from the Sierra Nevada, California. *Contributions to Mineralogy and Petrology* 151, 222–242.
- Lima, S.M., Corfu, F., Neiva, A.M.R., Ramos, M.F., 2011. Dissecting complex magmatic processes: an in-depth U–Pb study of the Pavia pluton, Ossa-Morena zone, Portugal. *Journal of Petrology* 53, 1887–1911.
- Linnemann, U., Gerdes, A., Drost, K., Buschmann, B., 2007. The continuum between Cadomian orogenesis and opening of the Rheic Ocean: constraints from LA-ICP-MS U–Pb zircon dating and analysis of plate-tectonic setting (Saxo-Thuringian zone, NE Bohemian massif, Germany). In: Linnemann, U., Nance, R.D., Kraft, P., Zulauf, G. (Eds.), *The Evolution of the Rheic Ocean: From Avalonian-Cadomian Active Margin to Alleghenian-Variscan Collision*. Geological Society of America Special Paper 423, 61–96.
- López-Plaza, M., López-Moro, F.J., 2003. Eurogranites Field Meeting. *The Tormes Dome: A Guide Book*, vol. 3, p. 192.
- Martínez Catalán, J.R., 2011. The Central Iberian arc: implications for the Iberian Massif. *Geogaceta* 50, 7–10.
- Martínez Catalán, J.R., Arenas, R., Abati, J., Sanchez Martinez, S., Diaz Garcia, F., Fernandez-Suarez, J., Gonzalez Cuadra, P., Castineiras, P., Gomez Barreiro, J., Diez Montes, A., Clavijo, E., Pascual, F.J., Andonaegui, P., Jeffries, T.E., Alcock, J.E., Diez Fernandez, R., Lopez Carmona, A., 2009. A rootless suture and the loss of the roots of a mountain chain: the Variscan Belt of NW Iberia. *Comptes Rendus Geoscience* 341, 114–126.
- Martínez-Catalán, J.R., Arenas, R., Diaz Garcia, F., Gomez-Barrero, J., Gonzalez Cuadra, P., Abati, J., Castineiras, P., Fernandez-Suarez, J., Sanchez Martinez, S., Andanaegui, P., Gonzalez Clavijo, E., Diez Montes, A., Rubio Pascual, F.J., Valle Aguado, B., 2007. Space and time in the tectonic evolution of the northwestern Iberian Massif. Implications for the comprehension of the Variscan belt. In: Hatcher Jr., R.D., Carlson, M.P., McBride, J.H., Martinez-Catalan, J.R. (Eds.), *4-D Framework of Continental Crust*. Geological Society of America Memoir, 2007.
- Martínez Catalán, J.R., Fernández Suárez, J., Jenner, G.A., Belousova, E., Diez Montes, A., 2004. Provenance constraints from detrital zircon U–Pb ages in the northwestern Iberian Massif: implications for Paleozoic plate configuration and Variscan evolution. *Journal of the Geological Society of London* 161, 461–473.
- Martins, H.C.B., Sant’Ovaia, H., Noronha, F., 2009. Genesis and emplacement of felsic Variscan plutons within a deep crustal lineation, the Penacova-Régua-Verín fault: an integrated geophysics and geochemical study (NW Iberian Peninsula). *Lithos* 11 (3–4), 142–155.
- Montero, P., Bea, F., Zinger, T.F., Scarrow, J.H., Molina, J.F., Whitehouse, M., 2004. 55 million years of continuous antesis in Central Iberia: single-zircon dating of the Peña Negra Complex. *Journal of the Geological Society of London* 161, 255–263.
- Moita, P., Santos, J.F., Pereira, M.F., 2009. Layered granitoids: interaction between continental crust recycling processes and mantle-derived magmatism. Examples from the Évora Massif (Ossa–Morena Zone, southwest Iberia, Portugal). *Lithos* 111, 125–141.
- Murphy, J.B., 2013. Appinite suites: a record of the role of water in the genesis, transport, emplacement and crystallization of magma. *Earth Science Reviews* 119, 35–59.
- Nance, R.D., Gutiérrez-Alonso, G., Keppie, J.D., Linnemann, U., Murphy, J.B., Quesada, C., Strachan, R.A., Woodcock, N.H., 2010. Evolution of the Rheic Ocean. *Gondwana Research* 17, 194–222.
- Neiva, A.M.R., Williams, I.S., Lima, S.M., Teixeira, R.J.S., 2012. U–Pb and ³⁹Ar/⁴⁰Ar data constraining the ages of the source, emplacement and recrystallization/cooling events from late- to post-D3 Variscan granites of the Gouveia area, central Portugal. *Lithos* 153, 72–83.
- Neiva, A.M.R., Williams, I.S., Ramos, J.M.F., Gomes, M.E.P., Silva, M.M.V.G., Antunes, I.M.H.R., 2009. Geochemical and isotopic constraints on the petrogenesis of Early Ordovician granodiorite and Variscan two-mica granites from the Gouveia area, central Portugal. *Lithos* 111, 186–202.
- Orejana, D., Villaseca, C., Pérez-Soba, C., López-García, J.A., Billström, K., 2009. The Variscan gabbros from the Spanish Central System: a case for crustal recycling in the sub-continental lithospheric mantle? *Lithos* 110, 262–276.
- Pankhurst, R.J., Weaver, S.D., Herve, F., Larrondo, P., 1999. Mesozoic-Cenozoic evolution of the North Patagonian batholith in Aysen, southern Chile. *Journal of the Geological Society* 156, 673–694.
- Parada, M.A., Nyström, J.O., Levi, B., 1999. Multiple sources for the Coastal Batholith of central Chile 31–348S: geochemical and Sr–Nd isotopic evidence and tectonic implications. *Lithos* 46, 504–521.
- Petford, N., Gallagher, K., 2001. Partial melting of mafic (amphibolitic) lower crust by periodic influx of basaltic magma. *Earth and Planetary Science Letters* 193, 483–489.
- Petford, N., Atherton, M., 1996. Na-rich partial melts from newly underplated basaltic crust: the Cordillera Blanca Batholith, Peru. *Journal of Petrology* 37, 1491–1521.
- Pereira, M.F., Castro, A., Chichorro, C., Fernández, C., Díaz-Alvarado, J., Martí, J., Rodríguez, C., 2014. Chronological link between deep-seated processes in magma chambers and eruptions: Permo-Carboniferous magmatism in the core of Pangaea (Southern Pyrenees). *Gondwana Research* 25, 290–308.
- Pereira, M.F., Chichorro, M., Silva, J.B., Ordoñez-Casado, B., Lee, J.K.W., Williams, I.S., 2012a. Early Carboniferous wrenching, exhumation of high-grade metamorphic rocks and basin instability in SW Iberia: constraints derived from structural geology and U–Pb and ⁴⁰Ar–³⁹Ar geochronology. *Tectonophysics* 558–559, 28–44.
- Pereira, M.F., Chichorro, M., Johnston, S.T., Gutiérrez-Alonso, G., Silva, J.B., Linnemann, U., Hofmann, M., Drost, K., 2012b. The missing Rheic Ocean magmatic arcs: provenance analysis of Late Paleozoic sedimentary clastic rocks of SW Iberia. *Gondwana Research* 22 (3–4), 882–891.
- Pereira, M.F., Chichorro, M., Williams, I.S., Silva, J.B., Fernandez, C., Diaz-Azpiroz, M., Apraiz, A., Castro, A., 2009. Variscan intra-orogenic extensional tectonics in the Ossa-Morena Zone (Évora-Aracena-Lora del Rio metamorphic belt, SW Iberian Massif): SHRIMP zircon U–Th–Pb geochronology. In: Murphy, J.B., Keppie, J.D., Hynes, A.J. (Eds.), *Ancient Orogens and Modern Analogues*. Geological Society, London, Special Publications 327, pp. 215–237.
- Rubio Pascual, F.J., Arenas, R., Martínez Catalán, J.R., Rodríguez Fernández, L.R., Wijbrans, J.R., 2013. Thickening and exhumation of the Variscan roots in the Iberian Central System: tectonothermal processes and ⁴⁰Ar/³⁹Ar ages. *Tectonophysics* 587, 207–221.
- Scarrow, J.H., Molina, J.F., Bea, F., Montero, P., 2009. Within-plate calc-alkaline rocks: insights from alkaline mafic magma–peraluminous crustal melt hybrid appinites of the Central Iberian Variscan continental collision. *Lithos* 110, 50–64.
- Sisson, T.W., Ratajeski, K., Hankins, W.B., Glazner, A.F., 2005. Voluminous granitic magmas from common basaltic sources. *Contributions to Mineralogy and Petrology* 148, 635–661.
- Solá, A.R., Williams, I.S., Neiva, A.M.R., Ribeiro, M.L., 2009. U–Th–Pb SHRIMP ages and oxygen isotope composition of zircon from two contrasting late Variscan granitoids, Nisa-Albuquerque batholith, SW Iberian Massif: Petrologic and regional implications. *Lithos* 111, 156–167.
- Stern, C.R., 1991. Role of subduction erosion in the generation of Andean magmas. *Geology* 19, 78–81.
- Stampfli, G.M. (Ed.), 2001. *Geology of the Western Swiss Alps, A Guide Book: Mémoires de Géologie (Lausanne)*. Introduction: Geology of the Western Swiss Alps, vol. 36, p. 195, pp. 1–41.
- Stampfli, G.M., Hochard, C., Vêard, C., Wilhem, C., von Raumer, J., 2013. The formation of Pangea. *Tectonophysics* 593, 1–19.
- Stampfli, G.M., von Raumer, J., Wilhem, C., 2011. The distribution of Gondwana-derived terranes in the Early Paleozoic. In: Gutiérrez-Marco, J.C., Rábano, I., García-Bellido, D. (Eds.), *Ordovician of the World*, vol. 14. Instituto Geológico y Minero de España, Cuadernos del Museo Geominero, pp. 567–574.
- Stampfli, G.M., Kozur, H., 2006. Europe from the Variscan to the Alpine cycles. In: Gee, D.G., Stephenson, R. (Eds.), *European Lithosphere Dynamics*. Geological Society of London Memoir 32, 57–82.
- Stampfli, G.M., Borel, G.D., 2002. A plate tectonic model for the Paleozoic and Mesozoic constrained by dynamic plate boundaries and restored synthetic oceanic isochrones. *Earth and Planetary Science Letters* 196, 17–33.
- Stevens, G., Villaros, A., Moyon, J.-F., 2007. Selective peritectic garnet entrainment as the origin of geochemical diversity in S-type granites. *Geology* 35 (1), 9–12.

- Ugidos, J.M., Armenteros, I., Barba, P., Valladares, M.I., Colmenero, J.R., 1997. Geochemistry and petrology of recycled orogen-derived sediments: a case study from upper Precambrian siliciclastic rocks of the Central Iberian Zone, Iberian Massif, Spain. *Precambrian Research* 84, 163–180.
- Valle Aguado, B., Azevedo, M.R., Schaltegger, U., Martínez Catalán, J.R., Nolan, J., 2005. U–Pb zircon and monazite geochronology of Variscan magmatism related to synconvergence extension in Central Northern Portugal. *Lithos* 82, 169–184.
- Valverde-Vaquero, P., 1997. An integrated field, geochemical and U–Pb geochronological study of the southwest Hermitage Flexure (Newfoundland Appalachians, Canada) and the Sierra De Guadarrama (Iberian Massif, Central Spain): a contribution to the understanding of the geological evolution of circum-Atlantic Peri-Gondwana. Unpublished Doctoral (PhD) thesis. Memorial University of Newfoundland.
- Valverde-Vaquero, P., Bento dos Santos, T., Clavijo, E.G., Díez Montes, A., Ribeiro, M.L., Solá, A.R., Dias da Silva, Í., 2011. The Berlengas Archipelago granitoids within the frame of the Variscan Orogeny, W Portugal: new data and insights. VII Hutton Symposium on Granites and Related Rocks. Abstracts Book, Avila, Spain, 4–9 July 2011, pp. 131–132.
- Villaseca, C., Bellido, F., Pérez-Soba, C., Billström, K., 2009. Multiple crustal sources for post-tectonic I-type granites in the Hercynian Iberian belt. *Mineralogy and Petrology* 96, 197–211.
- Vogt, K., Gerya, T.V., Castro, A., 2012. Crustal growth at active continental margins: numerical modeling. *Physics of the Earth and Planetary Interiors* 192–193, 1–20.
- von Raumer, J.F., Bussy, F., Schaltegger, U., Schulz, B., Stampfl, G.M., 2012. Pre-Mesozoic Alpine basement – their place in the European Paleozoic framework. *GSA Bulletin* 125 (1–2), 89–108.
- Weil, A., Gutiérrez-Alonso, G., Johnston, S.T., Pastor-Galán, D., 2012. Kinematic constraints on buckling a lithospheric-scale orocline along the northern margin of Gondwana: a geologic synthesis. *Tectonophysics* 582, 25–49.
- Wu, F.Y., Lin, J.Q., Wilde, S.A., Zhang, X.O., Yang, J.H., 2005. Nature and significance of the Early Cretaceous giant igneous event in Eastern China. *Earth and Planetary Science Letters* 233, 103–119.
- Wu, F.Y., Sun, D.Y., Li, H.M., Jahn, B.M., Wilde, S.A., 2002. A-type granites in north-eastern China: age and geochemical constraints on their petrogenesis. *Chemical Geology* 187, 143–173.
- Zeck, H.P., Wingate, M.T.D., Pooley, G.D., 2007. Ion microprobe U–Pb zircon geochronology of a late tectonic granitic-gabbroic rock complex within the Hercynian Iberian belt. *Geological Magazine* 144, 157–177.
- Zhang, Z.-M., Dong, X., Xiang, H., Liou, J.G., Santosh, M., 2013. Building of the deep Gangdese Arc, South Tibet: Paleocene plutonism and granulite-facies Metamorphism. *Journal of Petrology* 54, 2547–2580.